



## Ethylene resistance in flowering ornamental plants - improvements and future perspectives

Olsen, Andreas; Lütken, Henrik Vlk; Hegelund, Josefine Nymark; Müller, Renate

*Published in:*  
Horticulture Research

*DOI:*  
[10.1038/hortres.2015.38](https://doi.org/10.1038/hortres.2015.38)

*Publication date:*  
2015

*Document version*  
Publisher's PDF, also known as Version of record

*Document license:*  
[CC BY-NC-ND](#)

*Citation for published version (APA):*  
Olsen, A., Lütken, H. V., Hegelund, J. N., & Müller, R. (2015). Ethylene resistance in flowering ornamental plants - improvements and future perspectives. *Horticulture Research*, 2, [15038].  
<https://doi.org/10.1038/hortres.2015.38>

## REVIEW ARTICLE

## Ethylene resistance in flowering ornamental plants – improvements and future perspectives

Andreas Olsen, Henrik Lütken, Josefine Nymark Hegelund and Renate Müller

Various strategies of plant breeding have been attempted in order to improve the ethylene resistance of flowering ornamental plants. These approaches span from conventional techniques such as simple cross-pollination to new breeding techniques which modify the plants genetically such as precise genome-editing. The main strategies target the ethylene pathway directly; others focus on changing the ethylene pathway indirectly via pathways that are known to be antagonistic to the ethylene pathway, e.g. increasing cytokinin levels. Many of the known elements of the ethylene pathway have been addressed experimentally with the aim of modulating the overall response of the plant to ethylene. Elements of the ethylene pathway that appear particularly promising in this respect include ethylene receptors as ETR1, and transcription factors such as EIN3. Both direct and indirect approaches seem to be successful, nevertheless, although genetic transformation using recombinant DNA has the ability to save much time in the breeding process, they are not readily used by breeders yet. This is primarily due to legislative issues, economic issues, difficulties of implementing this technology in some ornamental plants, as well as how these techniques are publically perceived, particularly in Europe. Recently, newer and more precise genome-editing techniques have become available and they are already being implemented in some crops. New breeding techniques may help change the current situation and pave the way toward a legal and public acceptance if products of these technologies are indistinguishable from plants obtained by conventional techniques.

*Horticulture Research* (2015) 2, 15038; doi:10.1038/hortres.2015.38; published online: 26 August 2015

## INTRODUCTION

The production of potted ornamental plants and cut flowers is a growing industry with annual turnover in the tens of billions of dollars,<sup>1</sup> with the European Union being the single largest producer and consumer.<sup>2</sup> Due to the competitive nature of the ornamental plant industry, research and development of new products and improvement of quality in existing products have become essential. The quality of flowering ornamentals, often judged by the longevity of their flowers, is an extremely important parameter in assessing quality.<sup>3</sup> During the postharvest period, plants will likely experience stress as a result of poor lighting, temperature, and suboptimal humidity or watering.<sup>4,5</sup> This stress often leads to visual symptoms such as wilting, color change, and abscission of various plant parts including flowers, petals, and buds. In climacteric plants, stress triggers the production of the phytohormone ethylene, which quickly deteriorates plants visually. Ethylene of exogenous origin can also affect the plants in closed spaces<sup>6</sup> and plants that have been exposed to ethylene will often no longer be sellable.


In recent years, the biological significance of ethylene in ornamental production, its signaling pathway within the plant, and methods to alleviate its consequence to the aesthetic value of ornamental plants have been extensively reviewed.<sup>7–13</sup> However, many reviews only consider a limited number of plant species, or focus on a particular perspective for solving the problem in only one or two of the levels of the ethylene pathway. In order to acquire an overview of the problems associated with ethylene-induced quality losses in ornamental plant production, recent research and advances in the field of ethylene biology and breeding techniques must be considered. The present review aims to compare results from past approaches in order to discuss future breeding strategies.

It will also point out avenues not yet explored in the area of ornamental plant breeding which may be essential for reducing ethylene responses that remains a decisive factor for high-quality production of many ornamental plants.

## REGULATION OF THE ETHYLENE PATHWAY

The ethylene biosynthesis and signaling pathway can be presented in a linear model (Figure 1). Ethylene is essential for many processes in the plant and thus, there is a constant, low ethylene production (Figure 1a).<sup>14</sup> Under certain conditions, however, ethylene biosynthesis and sensitivity increase in specific tissues and this triggers the ethylene signaling pathway (Figure 1b).<sup>15</sup> This initially starts as an increase in expression of some of the enzymes responsible for ethylene biosynthesis<sup>16–18</sup> which leads to higher ethylene production<sup>19</sup> that may amplify itself in an autocatalytic fashion in some cases.<sup>20</sup>

Recently, it was shown that different ACS homologs of *Dianthus caryophyllus*<sup>18</sup> and *Dianthus superbus*<sup>21</sup> were expressed in association with either the basal or the climacteric phase. In *Petunia*<sup>22</sup> and *Paeonia suffruticosa*,<sup>23</sup> the same was observed for certain ACS and ACO homologs. It therefore seems that these enzymes, central for ethylene biosynthesis (Figure 1), may hold the key for the ability of the plant to transition from basal to climacteric ethylene production (Figure 1).<sup>24</sup> Furthermore in *Arabidopsis*, ACS activity was demonstrated to be affected by the formation of hetero- and homo-dimers,<sup>24,25</sup> and phosphorylation by the mitogen-activated protein kinase (MAPK) MPK6, which itself is expressed in response to stress.<sup>26</sup> Recently, a similar role for MPK6 was demonstrated in *Rosa hybrida*.<sup>27</sup>



**Figure 1.** Simplified ethylene pathway. **(a)** Basal production of ethylene in the flowers during development before senescence. **(b)** The ethylene pathway upon triggering. The stimulus is translated to elevated ethylene synthesis producing higher levels of ethylene which inactivates the receptors initiating the signaling cascade which changes gene expression and finally induces physiological processes in the flower which may include the initiation of an autocatalytic loop.<sup>14</sup> **(c)** Simplified molecular events of the ethylene pathway. Methionine is enzymatically converted to S-adenosyl-L-methionine (SAM) by SAM synthase (SAS). SAM is partially converted back to methionine via several steps, but it also produces 1-aminocyclopropane-1-carboxylic acid (ACC) by ACC synthase (ACS). ACC is transformed to ethylene by ACC oxidase (ACO). Ethylene binds to receptors and stops their signal to CONSTITUTIVE TRIPLE RESPONSE1 (CTR1), which then stops its suppressing signal to ETHYLENE INSENSITIVE2 (EIN2). The released EIN2 is then cleaved and part of it is transported into the nucleus where activation of the ETHYLENE INSENSITIVE3/ETHYLENE INSENSITIVE3-LIKE (EIN3/EIL) transcription factor family occurs. This initiates a transcription cascade by activation of ETHYLENE RESPONSE FACTORS (ERFs) which eventually leads to differential gene expression and a physiological response.<sup>15</sup>

The function of ethylene receptors too, is subject to modification from various proteins such as RESPONSIVE TO ANTAGONIST1 (RAN1),<sup>28,29</sup> REVERSION TO ETHYLENE SENSITIVITY1 (RTE1),<sup>30–32</sup> and their own histidine autokinase activity.<sup>33</sup> Receptors often act simultaneously, and in physical association with each other,<sup>34,35</sup> but as with the ethylene biosynthesis enzymes, different receptor homologs are expressed depending on the developmental stage of the flower, as has been documented in *D. caryophyllus*,<sup>36</sup> *R. hybrida*,<sup>37</sup> *Oncidium*,<sup>38</sup> *Delphinium*,<sup>39</sup> and *Pelargonium hortorum*.<sup>40</sup>

The function of EIN2 can be inhibited not only by CTR1 but also ETHYLENE RESISTANT/ETHYLENE RECEPTOR1 (ETR1)<sup>41</sup> and EIN2 TARGETING PROTEIN1 (ETP1) and ETP2. ETP1 and ETP2 were down-regulated in the presence of ethylene,<sup>42,43</sup> but activated CTR1 in the absence of ethylene.<sup>44</sup> Unlike most of the other genes of the ethylene pathway, EIN2 has no homologs and no functional overlap with other genes.<sup>45–47</sup> There is some indication that EIN2 expression is influenced by auxin and abscisic acid (ABA),<sup>48</sup> which means that EIN2 could be a point of crosstalk for several different pathways. Interestingly, increase in EIN2 protein levels has been shown to be concomitant with expression of the transcription factor ORESARA1 (ORE1) in *Arabidopsis*, associated with age-induced programmed cell death. ORE1 is suppressed by miRNA164, which declines with cell age.<sup>49</sup> In *R. hybrida* flowers exposed to ethylene, different miRNAs including miRNA164, exhibited a change in expression level in petals,<sup>50</sup> indicating a possible new level for expression control of ethylene genes.

EIN3/EIL homologs also change in the transition of the plant tissue to the climacteric phase. In *D. caryophyllus*, DcEIL3 increased<sup>51</sup> and DcEIL1 decreased<sup>52</sup> in expression as flower development occurred. Tanase *et al.*<sup>53</sup> on the other hand documented that DcEIL3 expression did not change with age in petals, but DcEIL4

was expressed less in older flowers. In *P. suffruticosa*, PsEIL1 accumulated from the flower opening stage to senescence, while PsEIL2 and PsEIL3 decreased after flower opening.<sup>54</sup> PsEIL2 and PsEIL3 mRNA levels increased in response to exogenous ethylene, while PsEIL1 was unaffected by this treatment and plants treated with 1-methylcyclopropane (1-MCP) and exposed to ethylene, exhibited a decrease in PsEIL3 expression.<sup>54</sup> In *Oncidium gardneri*, OgEIL1 and OgEIL2 were constitutively expressed in flower buds, but when the buds were exposed to ethylene, OgEIL1 clearly peaked in expression relative to OgEIL2.<sup>55</sup> Collectively, these genes present numerous targets for modifying ethylene regulation molecularly.

## REDUCING ETHYLENE-PROMOTED SENESCENCE

Approaches addressing the problem of ethylene-induced senescence can be broadly divided into two main groups: interventions directly addressing the ethylene pathway and those indirectly targeting it as described below. Because the ethylene pathway is fundamentally integrated into plant metabolism, changing a seemingly unrelated pathway of the plant can result in some effect on the ethylene pathway. Techniques to combat senescence such as changes in the gas composition of the atmosphere or temperature of storage places for plants have also been examined and gave mixed results.<sup>56–58</sup> These techniques, although successful to a point, lead to unsustainable production of ornamental plants because they extend the time needed for production and cost in terms of consumption of electricity, water, personnel, and specialized equipment. Silver thiosulfate (STS) and 1-MCP have been proven to be powerful inhibitors of ethylene responses and STS is commonly used in ornamental plant production. However, silver is harmful to humans and the environment and thus its use should be

avoided. Plant breeding on the other hand strives to improve the intrinsic quality of ornamental plants and thereby produce sustainable products.<sup>59</sup>

### TARGETING THE ETHYLENE PATHWAY DIRECTLY

Cross-pollination coupled with ethylene screening

The simple strategy of choosing individuals in a population that display superior flower ethylene tolerance and crossing these individuals with each other, will result in progeny with a lower ethylene sensitivity than the original population, if the trait is heritable.<sup>60</sup> Success in such breeding has been reported in *Begonia*<sup>61</sup> and *D. caryophyllus*.<sup>62</sup> It has long been known that there are vast differences among cultivars of *D. caryophyllus* in their ethylene biosynthetic ability and affinities of receptors to ethylene.<sup>63</sup> By using only simple pollination, Onozaki and coworkers<sup>62</sup> achieved a significant improvement in the longevity of *D. caryophyllus* flowers from 1992 to 2004. Their original material had flower longevity of 5.7 days,<sup>64,65</sup> but through repeated crossing, the sixth generation had increased its mean of flower life to 15.9 days.<sup>62</sup> An investigation into the cause of this increase, revealed that the ethylene biosynthesis enzyme genes *DcACS1*, *DcACS2*, and *DcACO1* were all expressed in extremely low levels in cultivars with a good longevity and ethylene production was as low.<sup>66,67</sup>

This approach involves screening for ethylene insensitivity and will only be applicable if a population exhibits significant differences in response between individuals. The number of ornamental flower cultures showing such diversity is, however, very large. Differences in longevity of flowers have been noted for *Paonia*,<sup>68</sup> *Delospermum*, *Campanula*, *Sedum*, *Cephalaria*, *Lobelia*, *Armeria*, *Primula*, *Penstemon*,<sup>69</sup> *Lisianthus*, *Trachelium*, *Zinnia*,<sup>70</sup> *Potentilla*, *Lysimachia*, *Veronica*, *Chelone*,<sup>71</sup> and *Rosa*<sup>11,72,73</sup> which also exhibited varying ethylene production levels.<sup>73–76</sup> Similar observations has been made for *Pelargonium*,<sup>77</sup> where heritability has also been documented,<sup>78</sup> as it has been for *Impatiens walleriana*,<sup>79</sup> *Antirrhinum majus*,<sup>80–82</sup> *Dianthus barbatus*,<sup>83</sup> and *Petunia*.<sup>84</sup> In many ornamental plant genera, cultivar-specific variation in ethylene sensitivity has been demonstrated including *Phalaenopsis*<sup>85,86</sup> and *Kalanchoe*.<sup>87</sup>

### Hybridization

Many of the abovementioned genera are typically represented by hybrids in ornamental plant production. Hybridization is achieved by interspecific cross-pollination and it thus produces extremely heterogeneous progeny, which may be good for producing cultivars with higher ethylene tolerance. A well-documented example of this is found in Australian waxflower breeding where *Chamaelium* species were hybridized with *Verticordia plumosa*, forming more ethylene-insensitive plants.<sup>88</sup> Hybrids of *Leptospermum* species<sup>89</sup> and *Grevillea*<sup>90</sup> similarly have been reported to have longer longevity than the parental species, which may be correlated with higher ethylene tolerance as both genera are climacteric. This aspect, however, has not yet been investigated.

The major disadvantage of both intra- and interspecific cross-pollination is that it cannot be used directly for improving existing cultivars. Considerable back-crossing to the original plant may be necessary.<sup>91</sup> This problem can be solved to some extent by employing marker-assisted selection, if reliable markers are found<sup>92</sup> and have been used successfully in ornamentals such as *D. caryophyllus* with regards to bacterial wilt resistance.<sup>93</sup> Hybridization can be more demanding as manipulation of the style as well as embryo rescue or ovule and ovary culture may be necessary in order to yield any progeny, as exemplified in *Kalanchoe* species hybridization.<sup>94</sup>

### Conventional mutagenesis

Techniques that increase genetic variability where no natural variation exists are found in conventional breeding with the use of

mutagenic chemicals or radiation.<sup>95</sup> Such breeding strives to change the cultivar in a certain qualitative aspect and leaves its agronomical traits unchanged, making it easier for producers to handle, and saves time that would otherwise be spent on back-crossing.<sup>71,96</sup> However, as it is impossible to control where the mutation occurs in the genome, many plants will have to be screened before a plant exhibiting any change in the relevant trait is found.<sup>97</sup> Targeting Induced Local Lesions In Genomes (TILLING) can be used to save time that would otherwise be needed for phenotyping the plants and has been used successfully in the field of ethylene-response improvement. Dahmani-Mardas *et al.*<sup>98</sup> produced *Cucumis melo* with a knockout mutation in the *CmACO1* gene which displayed a longer shelf life and better firmness than non-mutated plants. TILLING is readily applicable to ornamental plants as is exemplified in *Petunia*.<sup>99</sup>

### Genetic transformation

Genetic transformation is the transfer of foreign or native genes and promoters to a target genome by *Agrobacterium*-mediated transformation, particle bombardment, or infiltration. *Agrobacterium tumefaciens*-mediated transformation with recombinant DNA has been used in various ways to modify ethylene biosynthesis and signaling. These approaches have improved understanding of ethylene biosynthesis and signaling; however, their use in commercial breeding is limited and this technique's products are considered genetically modified.<sup>59,100</sup>

Antisense or sense transformations of ACS and ACO genes have been successfully attempted in various ornamental species including *D. caryophyllus*,<sup>101–113</sup> *Torenia*,<sup>104</sup> *Petunia*,<sup>22,105</sup> and *Begonia*.<sup>60,106</sup> All transformed plants exhibited longer shelf life, presumably due to lower production of ethylene. Klee *et al.*<sup>107</sup> conducted an alternative modification of the biosynthesis pathway with the removal of ACC by the addition of the enzyme ACC deaminase, found in bacteria, to *Solanum lycopersicum* which significantly delayed ripening of the fruit. However, even though plants with lower ethylene production have a longer flower life, their quality can still be negatively affected by exogenous ethylene.

Transformations of *Petunia* with the *Atert1-1* mutated gene from *Arabidopsis* using CaMV 35S constitutive promoter, resulted in plants with considerably higher ethylene tolerance but also with severely hampered growth.<sup>108</sup> Succeeding studies used a flower-specific FBP1 promoter in *Kalanchoe*,<sup>109</sup> *Campanula*,<sup>110,111</sup> and *D. caryophyllus*,<sup>112,113</sup> which provided a better flower longevity without other developmental effects, since the promoter ensured expression solely in flowers. Transformations of mutated genes other than *Atert1-1* have also been studied in *Nemesia strumosa*<sup>114</sup> and *Torenia fournieri*. In both cases, ethylene insensitivity was increased, but not as much as with *Atert1-1* (Table 1).<sup>8</sup>

Other promoters such as FBP3,<sup>115</sup> P<sub>SAG-12</sub>,<sup>116</sup> FS19, and FS26<sup>8</sup> that are solely associated with flower tissue, also have potential to be used in this context and other promoters are still being investigated.<sup>117</sup> Chemically inducible promoter systems are also within the reach of today's technology, and are exemplified by the DEX-inducible system, demonstrated in *Petunia hybrida*<sup>118</sup> (Table 1), and more famously, the ethanol-inducible system<sup>119</sup> which has been successfully used in different crops.<sup>120</sup> Inducible systems seem not to affect the plant adversely,<sup>121</sup> but the inducing chemical still needs to be applied to the plant in due time, which makes such a system commercially less appealing.

Great care needs to be exercised when choosing a promoter, as a study by Cobb *et al.*<sup>122</sup> demonstrated that *Petunia* transformed with *Atert1-1* did not have delay in senescence under the flower-specific APETALA3 (AP3) promoter, as this promoter only drives expression in buds and young flowers, but not in mature flowers (Table 1).

Overexpressing *PhEIN2* in *Petunia* using the CaMV 35S promoter has also been attempted. This produced plants with significant



**Table 1.** Transformation of ornamental plants modifying ethylene receptors

Promoter	Gene	Plant species	<i>Agrobacterium tumefaciens</i> strain	Plasmid	Reference
CaMV 35S	<i>Atetr1-1</i>	<i>P. hybrida</i>	ABI	pMON11063	108
<i>Petunia</i> FBP1 CaMV 35S	<i>Atetr1-1</i>	<i>D. caryophyllus</i>	AGLO	pBEO210	
				pBEO220	112
AP3	<i>Atetr1-1</i>	<i>P. hybrida</i>	?	?	116
CaMV 35S	<i>Brassica oleracea</i> ers (boers)	<i>P. hybrida</i>	LBA4404	pBOERS4421	123
CaMV 35S	<i>CmETR1/H69A</i>	<i>N. strumosa</i>	GV2260	pBICm-ETR1/ H69A	113
CaMV 35S	<i>CmERS1/H70A</i>	<i>Lotus japonicus</i>	<i>Mesorhizobium loti</i> , MAFF303099	pGD499	124
<i>Petunia</i> FBP1	<i>Atetr1-1</i>	<i>Kalanchoe blossfeldiana</i>	AGLO	pBEO210	109
<i>Petunia</i> FBP1	<i>Atetr1-1</i>	<i>Oncidium, Odontoglossum</i>	EHA105 and LBA4404	pBEO210	125
<i>Petunia</i> FBP1	<i>Atetr1-1</i>	<i>Campanula carpatica</i>	AGLO	pBEO210	110
CaMV 35S	<i>DcETR1nr</i>	<i>T. fournieri</i>	?	pBIDc-ETR1nr	53
GVG DEX-inducible	<i>Atetr1-1</i>	<i>P. hybrida</i>	LBA4404	pTA7001	118

delays in petal senescence in response to exogenous ethylene or pollination, as well as inhibited root development and a shorter life span.<sup>126</sup> When this silencing transformation was combined with the *Atetr1-1* mutation, ethylene was inhibited even more,<sup>127</sup> demonstrating the possibility of increasing insensitivity by stacking transformations of several components simultaneously.

*Petunia* plants transformed with a sense *PhEIN3* exhibited delayed petal senescence, but no increase of flower longevity,<sup>8,127</sup> which may possibly be due to the redundancy found among EIN3/EIL members. *S. lycopersicum* was transformed with a construct silencing EIN3 BINDING F-BOX1 (EBF1) and EBF2, which rapidly target EIN3/EILs for degradation that significantly increased the rate of senescence in *S. lycopersicum*.<sup>45</sup> This presents an opportunity for future research, potentially by overexpression of EBFs.

ERFs are a sizable and diverse group of genes, many of which are up-regulated in response to ethylene, but have largely been overlooked by researchers so far. Chang et al.<sup>128</sup> observed that by silencing the ERF *Petunia* transcription factor homeodomain-leucine zipper protein (PhHDZip), *PhACS* and *PhACO* were decreased in expression in transformed flowers and led to an increase of flower longevity by 20%.

#### Genome-editing technologies

During the last decades, several innovative biotechnologies have been developed which appear to have potential in breeding toward ethylene tolerance.<sup>100,129,130</sup> These technologies are based on engineered nucleases that cleave DNA in a sequence-specific manner, thus enabling targeted genome-editing. Modifying or inactivating specific gene function is possible due to sequence-specific DNA binding domains or RNA sequences. The oldest of these techniques is zinc finger nucleases (ZNFs), which relies on an engineered endonucleases that are able to attach to a specific target sequence and induce a double-strand break. This is perceived by the cell, which repairs the break by mechanisms which may cause sequence alterations or introduce small templates.<sup>131,132</sup> A newer alternative to ZNFs is transcription activator-like effector nucleases (TALENs), which functions in much the same manner as ZNFs. Recently, however, the clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated protein9 (CRISPR/Cas9) system has been developed. The CRISPR/Cas9 system functions via a mechanism similar to RNA interference, which can recognize and cleave foreign DNA.<sup>100</sup> Both TALEN and CRISPR/Cas9 have been demonstrated to be functional for targeted mutagenesis in *S. lycopersicum*,<sup>133,134</sup> however, no modifications of the ethylene pathway have yet been conducted. It is important to mention that all these methods rely on a technology in order to deliver the construct to the genome of the plant, and are therefore still limited to plants where a genetic transformation, regeneration or

virus-based delivery system is reliable. In order to bypass negative effects on growth of plants due to defective ethylene signaling, the genes that should be targeted are those which have homologs that are clearly associated with the climacteric phase. Using this approach, the pathway should function normally under growth and development. Therefore, certain homologs of ACS, ACO, ethylene receptors, and EIN3/EIL may present good targets for knockouts.

#### TARGETING THE ETHYLENE PATHWAY INDIRECTLY

##### Hormonal interaction

Application of cytokinins such as kinetin and zeatin to petals of *D. caryophyllus* delayed the conversion of ACC to ethylene.<sup>135</sup> Cytokinin oxidase/dehydrogenase, responsible for cytokinin degradation, was up-regulated during senescence in *D. caryophyllus*<sup>136</sup> and *Petunia*,<sup>137</sup> and when it was inhibited in *D. caryophyllus* petals, the senescence phase was prevented.<sup>138</sup> This knowledge has been applied by transformations with isopentenyl transferase, important for the synthesis of many cytokinins, under the control of senescence-associated promoter P<sub>SAG12</sub>. Transformed *Petunia* plants exhibited elevated levels of cytokinins in flowers and significant delay in senescence and ethylene production as well as higher tolerance for exogenous ethylene.<sup>139</sup> Similarly in *Rosa*, ethylene sensitivity decreased in leaves, but flowers were not studied.<sup>140</sup>

Exogenous application of hormones and other substances have also been shown to decrease the expression of genes of the ethylene pathway including ABA, which inhibited ACS and ACO in *Hibiscus*<sup>141</sup> and *D. caryophyllus*.<sup>142</sup> Moreover, nitric oxide down-regulated the activity of *RhACO* and lowered the production of ethylene resulting in longer shelf life of *R. hybrida* flowers<sup>143</sup> and glucose down-regulated *PsACS1* in *P. suffruticosa* flowers.<sup>144</sup> Other sugars have also been shown to increase flower longevity.<sup>136,145–148</sup> These studies demonstrate that there is a vast potential in exploring new ways to achieve products of higher quality.

##### Senescence-related and non-ethylene pathway genes

Exploration of different genes which is seemingly not connected to the ethylene pathway has also been pursued. There are three main strategies in this area. The first is identification of genes that are highly expressed in young tissue but not during senescence, and constitutively overexpressing those. The second approach goes the other way and starts with the identification of genes highly expressed in senescing tissue, and silencing their expression. The third tactic is targeting protein synthesis.

The first strategy can be exemplified by the gene *FOREVER YOUNG FLOWER* (FYF) from *Arabidopsis*, which was highly expressed in young flowers but not in old flowers. Transformation using this gene under constitutive expression in *S. lycopersicum* resulted in

down-regulation of various ACS and ACO homologs. The same transformation in *Eustoma grandiflorum* caused a delay in senescence and down-regulation of *ERFs*.<sup>149</sup> Furthermore, when combined with *etr1*, *ein2*, or *ctr1* mutations, it further enhanced flower longevity in *Arabidopsis*.<sup>149</sup>

Investigating a MADS-domain transcriptional regulator, AGAMOUS-LIKE-15 (AGL15), Fernandez *et al.*<sup>150</sup> noted its accumulation in young tissue and developing organs. Constitutive overexpression of *AGL15* increased the longevity of petals significantly in *Arabidopsis*<sup>151</sup> and caused a repression of *GmACO1* in *Glycine max*.<sup>152</sup> It is now established that AGL15 has similar activity as AGL18, both serving as repressors of early flowering and in this way leading to better longevity if still expressed when flowering does occur.<sup>153</sup> Another MADS-domain transcriptional regulator that has been recognized to be associated with young flowers of *Phalaenopsis equestris* is PeMADS6. *Arabidopsis* plants transformed with a construct overexpressing *PeMADS6* exhibited flower longevity up to four times longer than wild type plants.<sup>154</sup>

Examples from the second strategy of silencing senescence-related genes, include *ACTIN-RELATED PROTEIN 4* (*ARP4*), a chromatin modification gene, that has been silenced using RNAi in *Arabidopsis*,<sup>155</sup> which resulted in longer flower life. The gene *MjXB3*, coding for a RING zinc finger ankyrin repeat protein, has been demonstrated to be highly expressed in senescing flowers of *Mirabilis jalapa*, *P. hybrida*, and *D. caryophyllus*. Using virus-induced gene silencing for *MjXB3* in *P. hybrida*, Xu *et al.*<sup>156</sup> demonstrated that flower longevity was extended by two days, corresponding to a 20% increase compared to wild type flowers.

The third strategy of targeting protein synthesis owes its inspiration to ethylene-insensitive flowers,<sup>13</sup> where complete inhibition of protein synthesis in flowers increases flower longevity.<sup>157</sup> The relevance of this to climacteric flowers has been attested by silencing *PBB2* (coding for the beta subunit of the 26S proteasome) using an inducible system in *Petunia*.<sup>158</sup> Mature cut flowers that were induced lasted considerably longer than uninduced flowers. Reid and Jiang<sup>13</sup> demonstrated the same concept by silencing one of the ribosomal subunit genes (*RPL2*), again using an inducible system. Once more, the longevity of cut *Petunia* flowers placed in water with the inducing agent was much better than those uninduced. This approach, however, still relies on the application of an inducer, and is not realistic for use in commercial potted-plant production. For the cut-flower industry, however, there may be great potential.

#### Natural transformation

Transformation with the naturally occurring soil bacterium *Agrobacterium rhizogenes* has been termed natural transformation<sup>159</sup> due to the fact that no recombinant DNA is used and the infection is a natural process. Authorities in Denmark have confirmed that naturally transformed plants are not covered by the GMO legislation in Europe.<sup>59</sup> Using a wild strain of *A. rhizogenes*, Lütken *et al.*<sup>59,160</sup> transformed *Kalanchoë* plants and observed that plants which contained *rol*-genes from *A. rhizogenes* exhibited significantly increased postharvest quality,<sup>159</sup> an ability which was maintained through two generations along with the *rol*-genes.<sup>161</sup> The mechanism behind this response is yet unknown and many open reading frames of the plasmids in question still remain uninvestigated. Changes in longevity of flowers transformed with *rol*-genes may be due to altered hormone homeostasis and/or sugar metabolism and transport.<sup>159</sup>

## DISCUSSION

Strategies targeting the ethylene pathway indirectly seem to be as effective as those which target it directly. Of the indirect strategies, ectopic overexpression of genes associated with juvenile tissue, such as *FYF* or *AGL15*, seems to be particularly promising, as is the prospect of overproduction of hormones that are antagonistic to

ethylene, such as cytokinins and ABA. Strategies that focus on the ethylene pathway directly have been more numerous and have targeted nearly all known elements of the ethylene pathway. Research comprising flower-specific promoter sequences in genetic transformations seems to be especially propitious in providing the ability to express the transferred gene only in flowers or even more specifically, only in senescing flowers. Silencing several genes simultaneously in the flowers of plants as in *Petunia*,<sup>119</sup> may be the ultimate answer, and is a feature that is well within the grasp of current technology and methodology. Various combinations of overexpression of certain genes associated with development and silencing genes associated with senescence may very well be a worthwhile strategy as well.

It is remarkable that both conventional techniques such as simple cross-pollination and genetic transformation using recombinant DNA succeeded in breeding ornamental plants which have higher tolerance for ethylene and thus higher intrinsic quality. *Agrobacterium*-mediated transformation, where reliable, has the advantage that breeding can be much faster when transferring specific target genes. Breeding based on genetic transformation is more precise and can easily cross incompatibility boundaries in comparison with conventional crossing.<sup>91</sup> This is a particularly important advantage in ornamental plants that have long development cycles such as orchids,<sup>162</sup> however, in fast growing plants, this advantage is less substantial.<sup>60</sup>

Although conventional and genetic transformation are not mutually exclusive, the fact remains that ornamental plant products that are genetically modified are conspicuously missing on the market, especially in Europe, with only a handful of products sold worldwide.<sup>163</sup> The basic reason for this is the status of such product as genetically modified organisms (GMOs). Europe, the biggest consumer, producer and breeder of ornamentals has some of the most comprehensive and strict legislation concerning GMOs.<sup>1</sup> At the moment, the approval of GMO products is conducted on a case-by-case basis and is both expensive and time-consuming.<sup>164</sup> Moreover, the technology and training required for producing GMOs are themselves expensive, which brings a substantial economic burden.<sup>60</sup> Producers not operating in Europe also face difficulties as GMO legislation may be quite different from country to country.<sup>165</sup> However, the emergence of new genome-editing techniques presents new challenges particularly for European legislation in the area of GMOs, which could lead the way to broader acceptance of new breeding techniques and their products.

Techniques such as ZFN, TALENs, and CRISPR/Cas9 that can be used to achieve precise mutagenesis and silencing or overexpressing genes are now readily available, and can be used in such a way that does not introduce foreign elements to the genome of the final product.<sup>127,166</sup> The question of whether the products of these techniques are considered GMOs or not has not yet been settled, and jurisdictions that consider the methods alone rather than the actual properties of the final plant product may end up with ineffective legislation that cannot be enforced, as pointed out in *Seeds of Change*.<sup>167</sup> Certainly such techniques have the ability to greatly reduce the time, and thus the bulk of the cost which is associated with plant breeding. The first commercially available plant derived from a genome-editing technique, a herbicide-resistant oilseed rape, has already been announced.<sup>168</sup> There are still technical difficulties associated with the actual transformation step for many ornamental plants, which can only be overcome by further research. For this reason, it is must be of the utmost importance for any plant breeding company to devote some of its resources to research in such new breeding techniques.

## CONCLUSIONS

Strategies targeting the ethylene pathway directly seem to have as much potential as those targeting indirect components, and very

diverse species of ornamental plants suggest great potential in ethylene tolerance breeding. Although conventional breeding techniques are slower than newer breeding techniques, they remain the most used in ornamental plant breeding for higher ethylene tolerance. This is most probably due to the legal status of genetic modification approaches. However, as newer precise genome-editing techniques become available, it is very likely that the products of such techniques will be accepted worldwide. The reason for this is that such plants would be indistinguishable from plants derived by the use of conventional breeding techniques. It is therefore recommendable for ornamental plant breeders to begin the implementation of such new breeding technologies, as they have great potential to lead to superior plant products needed by the ornamental plant industry.

## COMPETING INTERESTS

The authors have no conflicts to declare.

## REFERENCES

- Chandler SF, Brugliera F. Genetic modification in floriculture. *Biotechnol Lett* 2011; **33**: 207–214.
- European Commission Agriculture and Rural Development. *Statistics for live plants and products of floriculture, 2013*. Available at [http://ec.europa.eu/agriculture/fruit-and-vegetables/product-reports/flowers/index\\_en.htm](http://ec.europa.eu/agriculture/fruit-and-vegetables/product-reports/flowers/index_en.htm) (accessed 1 March 2014).
- Palma M, Hall C, Collart A. Repeat buying behavior for ornamental plants: A consumer profile. *JFDRS* 2011; **42**: 67–77.
- Wagstaff C, Bramke I, Breeze E, Thornber S, Harrison E et al. A specific group of genes respond to cold dehydration stress in cut *Alstroemeria* flowers whereas ambient dehydration stress accelerates developmental senescence expression patterns. *J Exp Bot* 2010; **61**: 2905–2921.
- Müller R, Sisler EC, Serek M. Stress induced ethylene production, ethylene binding and the response to the ethylene action inhibitor (1-MCP) in miniature roses. *Sci Hortic* 1999; **83**: 51–59.
- Morgan PW. Another look at interpreting research to manage the effects of ethylene in ambient air. *Crop Sci* 2011; **51**: 903–913.
- Shibuya K. Molecular mechanisms of petal senescence in ornamental plants. *J Jpn Soc Hortic Sci* 2012; **81**: 140–149.
- Satoh S. Ethylene production and petal wilting during senescence of cut carnation (*Dianthus caryophyllus*) flowers and prolonging their vase life by genetic transformation. *J Jpn Soc Hortic Sci* 2011; **80**: 127–135.
- Fischer AM. The complex regulation of senescence. *Crit Rev Plant Sci* 2012; **31**: 124–147.
- Schaller GE. Ethylene and the regulation of plant development. *BMC Biol* 2012; **10**: 9.
- Fanourakis D, Pieruschka R, Savvides A, Macnisch AJ, Sarlikioti V et al. Sources of vase life variation in cut roses: A review. *Postharvest Biol Technol* 2013; **78**: 1–15.
- Merchante C, Alonso JM, Stepanova AN. Ethylene signaling: Simple ligand, complex regulation. *Curr Opin Plant Biol* 2013; **16**: 554–560.
- Reid MS, Jiang C. Postharvest biology and technology of cut flowers and potted plants. In: Janek J (ed.) *Horticultural reviews* 2012; Vol. 40. 1st ed. Hoboken, NJ: John Wiley & Sons, 1–54.
- Paul V, Pandey R, Srivastava GC. The fading distinctions between classical patterns of ripening in climacteric and non-climacteric fruit and the ubiquity of ethylene—an overview. *J Food Sci Technol* 2012; **49**: 1–21.
- Lin Z, Zhong S, Grierson D. Recent advances in ethylene research. *J Exp Bot* 2009; **60**: 3311–3336.
- Woodson WR, Park KY, Drory A, Larsen PB, Wang H. Expression of ethylene biosynthetic pathway transcripts in senescing carnation flowers. *Plant Physiol* 1992; **99**: 526–532.
- ten Have A, Woltering EJ. Ethylene biosynthetic genes are differentially expressed during carnation (*Dianthus caryophyllus* L.) flower senescence. *Plant Mol Biol* 1997; **34**: 89–97.
- In BC, Strable J, Binder BM, Falbel TG, Patterson SE. Morphological and molecular characterization of ethylene binding inhibition in carnations. *Postharvest Biol Technol* 2013; **86**: 272–279.
- Ma N, Tan H, Liu XH, Xue JQ, Li YH et al. Transcriptional regulation of ethylene receptor and CTR genes involved in ethylene-induced flower opening in cut rose (*Rosa hybrida*) cv. Samantha. *J Exp Bot* 2006; **57**: 2763–2773.
- Kende H. Ethylene biosynthesis. *Annu Rev Plant Physiol Plant Mol Biol* 1993; **44**: 283–307.
- Harada T, Murakoshi Y, Torii Y, Tanase K, Onozaki T et al. Analysis of genomic DNA of DCA1, a 1-aminocyclopropane-1-carboxylate synthase gene, expressed in senescing petals of carnation (*Dianthus caryophyllus*) and its orthologous genes in *D. superbus* var. *longicalycinus*. *Plant Cell* 2011; **30**: 519–527.
- Huang LC, Lai UL, Yang SF, Chu MJ, Kuo CI et al. Delayed flower senescence of *Petunia hybrida* plants transformed with antisense broccoli ACC synthase and ACC oxidase genes. *Postharvest Biol Technol* 2007; **46**: 47–53.
- Zhou L, Zhang C, Fu J, Liu M, Zhang Y et al. Molecular characterization and expression of ethylene biosynthetic genes during cut flower development in tree peony (*Paeonia suffruticosa*) in response to ethylene and functional analysis of PsACS1 in *Arabidopsis thaliana*. *J Plant Growth Regul* 2013; **32**: 362–375.
- Tsuchisaka A, Yu G, Jin H, Alonso JM, Ecker JR et al. A combinatorial interplay among the 1-Aminocyclopropane-1-Carboxylate Isoforms regulates ethylene biosynthesis in *Arabidopsis thaliana*. *Genetics* 2009; **183**: 979–1003.
- Wang KLC, Li H, Ecker JR. Ethylene biosynthesis and signaling networks. *Plant Cell* 2002; **14**: 131–151.
- Liu Y, Zhang S. Phosphorylation of 1-aminocyclopropane-1-carboxylic acid synthase by MPK6, a stress-responsive mitogen-activated protein kinase, induces ethylene biosynthesis in *Arabidopsis*. *Plant Cell* 2004; **16**: 3386–3399.
- Meng Y, Ma N, Zhang Q, You Q, Li N et al. Precise spatio-temporal modulation of ACC synthase by MPK6 cascade mediates the response of rose flowers to rehydration. *Plant J* 2014; **79**: 941–950.
- Hirayama T, Kieber JJ, Hirayama N, Kogan M, Guzman P et al. RESPONSIVE-TO-ANTAGONIST1, a Menkes/Wilson disease-related copper transporter, is required for ethylene signaling in *Arabidopsis*. *The Cell* 1999; **97**: 383–393.
- Binder BM, Rodríguez F, Bleeker AB. The copper transporter RAN1 is essential for biogenesis of ethylene receptors in *Arabidopsis*. *J Biol Chem* 2010; **285**: 37263–37270.
- Qiu L, Xie F, Yu J, Wen CK. *Arabidopsis* RTE1 is Essential to Ethylene Receptor ETR1 Amino-terminal signaling independent of CTR1. *Plant Physiol* 2012; **159**: 1263–1276.
- Yu Y, Wang H, Liu J, Fu Z, Wang J et al. Transcriptional regulation of two RTE-like genes of carnation during flower senescence and upon ethylene exposure, wounding treatment and sucrose supply. *Plant Biol* 2011; **13**: 719–724.
- Ma Q, Du W, Brandizzi F, Giovannoni JJ, Barry CS. Differential control of ethylene responses by GREEN-RIPE and GREEN-RIPE-LIKE1 provides evidence for distinct ethylene signaling modules in tomato. *Plant Physiol* 2012; **160**: 1968–1984.
- Hall BP, Shakeel SN, Amir M, Ul Haq N, Qu X et al. Histidine kinase activity of the ethylene receptor ETR1 facilitates the ethylene response in *Arabidopsis*. *Plant Physiol* 2012; **159**: 682–695.
- Grefen C, Städele K, Ruzicka K, Obrdlík P, Harter K et al. Subcellular localization and *in vivo* interactions of the *Arabidopsis thaliana* ethylene receptor family members. *Mol Plant* 2008; **1**: 308–320.
- Gao Z, Wen CK, Binder BM, Chen YF, Chang J et al. *Arabidopsis* receptors mediate signaling in heteromeric interactions among ethylene. *J Biol Chem* 2008; **283**: 23801–23810.
- Shibuya K, Nagata M, Tanikawa N, Yoshioka T, Hashiba T et al. Comparison of mRNA levels of three ethylene receptors in senescing flowers of carnation (*Dianthus caryophyllus* L.). *J Exp Bot* 2002; **53**: 399–406.
- Müller R, Stummann BM, Serek M. Characterization of an ethylene receptor family with differential expression in rose (*Rosa hybrida* L.) flowers. *Plant Cell Rep* 2000; **19**: 1232–1239.
- Huang WF, Huang PL, Do YY. Ethylene receptor transcript accumulation patterns during flower senescence in *Oncidium 'Gower Ramsey'* as affected by exogenous ethylene and pollinia cap dislodgement. *Postharvest Biol Technol* 2007; **44**: 97–94.
- Tanase K, Ichimura K. Expression of ethylene receptors DI-ERS1-3 and DI-ERS2, and ethylene response during flower senescence in *Delphinium*. *J Plant Physiol* 2006; **163**: 1159–1166.
- Dervinis C, Clark DG, Barrett JE, Nell TE. Effect of pollination and exogenous ethylene on accumulation of ETR1 homologue transcripts during flower petal abscission in geranium (*Pelargonium hortorum* L. H. Bailey). *Plant Mol Biol* 2000; **42**: 847–856.
- Bisson MMA, Groth G. New insight in ethylene signaling: autokinase activity of ETR1 modulates the interaction of receptors and EIN2. *Mol Plant* 2010; **5**: 882–889.
- Qiao H, Chang KN, Yazaki JH, Ecker JR. Interplay between ethylene, ETP1/ETP2 F-box proteins, and degradation of EIN2 triggers ethylene responses in *Arabidopsis*. *Genes Dev* 2009; **23**: 512–521.
- Bisson MMA, Groth G. Cyanide is an adequate agonist of the plant hormone ethylene for studying signaling of sensor kinase ETR1 at the molecular level. *Biochem J* 2012; **444**: 261–267.
- Shahri W, Tahir I. Flower senescence: Some molecular aspects. *Planta* 2014; **239**: 277–297.



- 45 Yang TF, Gonzalez-Carranza ZH, Maunders MJ, Roberts JEA. Ethylene and the regulation of senescence processes in transgenic *Nicotiana sylvestris* Plants. *Ann Bot* 2008; **101**: 301–310.
- 46 Yang Y, Wu Y, Pirrello J, Regad F, Bouzayen M et al. Silencing Sl-EBF1 and Sl-EBF2 expression causes constitutive ethylene response phenotype, accelerated plant senescence, and fruit ripening in tomato. *J Exp Bot* 2010; **61**: 697–708.
- 47 Fu Z, Wang H, Liu J, Liu J, Wang J et al. Cloning and characterization of a DCEIN2 gene responsive to ethylene and sucrose in cut flower carnation. *Plant Cell Tissue Organ Cult* 2011; **105**: 447–455.
- 48 Gao F, Hao J, Yao Y, Wang X, Hasi A. Cloning and characterization of ethylene-insensitive 2 (EIN2) Gene from *Cucumis melo*. *Rus J Plant Physiol* 2013; **60**: 713–719.
- 49 Kim JH, Woo HR, Kim J, Lim PO, Lee IC et al. Trifurcate feed-forward regulation of age-dependent cell death involving miR164 in Arabidopsis. *Science* 2009; **323**: 1053–1057.
- 50 Pei H, Ma N, Chen J, Zheng Y, Tian J et al. Integrative analysis of miRNA and mRNA profiles in response to ethylene in rose petals during flower opening. *PLoS One* 2013; **8**: e64290.
- 51 Iordachescu M, Verlinden S. Transcriptional regulation of three EIN3-like genes of carnation (*Dianthus caryophyllus* L. cv. Improved White Sim) during flower development and upon wounding, pollination, and ethylene exposure. *J Exp Bot* 2005; **1**: 2011–2018.
- 52 Waki K, Shibuya K, Yoshioka T, Hashiba T, Satoh S. Cloning of a cDNA encoding EIN3-like protein DC-EIL1 and decrease in its mRNA level during senescence in carnation flower tissues. *J Exp Bot* 2001; **52**: 377–379.
- 53 Tanase K, Aida R, Yamaguchi H, Tanikawa N, Nagata M et al. Heterologous expression of a mutated carnation ethylene receptor gene, Dc-ETR1, suppresses petal abscission and autocatalytic ethylene production in transgenic *Torenia fournieri* Lind. *J Jpn Soc Hortic Sci* 2011; **80**: 113–120.
- 54 Wang Y, Zhang C, Peiyi J, Wang X, Wang W et al. Isolation and expression analysis of three EIN3-like genes in tree peony (*Paeonia suffruticosa*). *Plant Cell* 2013; **112**: 181–190.
- 55 Chen SY, Tsai HC, Raghu R, Do YY, Huang PL. cDNA cloning and functional characterization of ETHYLENE INSENSITIVE3 orthologs from *Oncidium Gower Ramsey* involved in flower cutting and pollinia cap dislodgement. *Plant Physiol Biochem* 2011; **49**: 1209–1219.
- 56 Serek M, Woltering EJ, Sisler EC, Frello S, Sriskandarajah S. Controlling ethylene responses in flowers at the receptor level. *Biotechnol Adv* 2006; **24**: 368–381.
- 57 Yangkhamman P, Tanase K, Ichimura K, Fukai S. Depression of enzyme activities and gene expression of ACC synthase and ACC oxidase in cut carnation flowers under high-temperature conditions. *Plant Growth Regul* 2007; **53**: 155–162.
- 58 Kou L, Turner ER, Luo Y. Extending the shelf life of edible flowers with controlled release of 1-Methylcyclopropene and modified atmosphere packaging. *J Food Sci* 2012; **77**: 188–193.
- 59 Lütken H, Clarke JL, Müller R. Review: Genetic engineering and sustainable production of ornamentals: Current status and future directions. *Plant Cell Rep* 2012; **31**: 1141–1157.
- 60 Onozaki T. Improvement of flower vase life using cross-breeding techniques in carnation (*Dianthus caryophyllus* L.). *Jarg-Jpn Agr Res Q* 2008; **42**: 137–144.
- 61 Hvostlef-Eide AK, Fjeld T, Einset JW. Breeding Christmas begonia *Begonia × cheimanthus* Everett for increased keeping quality by traditional and biotechnological methods. *Acta Hortic* 1995; **405**: 197–204.
- 62 Onozaki T, Yagi M, Tanase K, Shibata M. Crossings and selections for six generations based on flower vase life to create lines with ethylene resistance or ultra-long vase life in carnations (*Dianthus caryophyllus* L.). *J Jpn Soc Hortic Sci* 2011; **80**: 486–498.
- 63 Wu MJ, Zacarias L, Reid MS. Variation in the senescence of carnation (*Dianthus caryophyllus* L.) cultivars. II. Comparison of sensitivity to exogenous ethylene and of ethylene binding. *Sci Hortic* 1991; **48**: 109–116.
- 64 Onozaki T, Ikeda H, Yamaguchi T. Genetic improvement of vase life of carnation flowers by crossing and selection. *Sci Hortic* 2001; **87**: 107–120.
- 65 Onozaki T, Ikeda H, Shibata M. Video evaluation of ethylene sensitivity after anthesis in carnation (*Dianthus caryophyllus* L.) flowers. *Sci Hortic* 2004; **99**: 187–197.
- 66 Tanase K, Onozaki T, Satoh S, Shibata M, Ichimura K. Differential expression levels of ethylene biosynthetic pathway genes during senescence of long-lived carnation cultivars. *Postharvest Biol Technol* 2008; **47**: 210–217.
- 67 Tanase K, Otsu S, Satoh S, Onozaki T. Expression and regulation of senescence-related genes in carnation flowers with low ethylene production during senescence. *J Jpn Soc Hortic Sci* 2013; **82**: 179–187.
- 68 Heuser CW, Evensen KB. Cut flower longevity of peony. *J Am Soc Hortic* 1986; **111**: 896–899.
- 69 van Doorn WG. Categories of petal senescence and abscission: A re-evaluation. *Ann Bot* 2001; **87**: 447–456.
- 70 Clark EMR, Dole JM, Carlson AS, Moody EP, McCall IF et al. Vase life of new cut flower cultivars. *Hort Technol* 2010; **20**: 1016–1025.
- 71 Woltering EJ, van Doorn WG. Role of ethylene in senescence of petals morphological and Taxonomical relationships. *J Exp Bot* 1988; **39**: 1605–1616.
- 72 Müller R, Andersen AS, Serek M. Differences in display life of miniature potted roses (*Rosa hybrida* L.). *Sci Hortic* 1998; **76**: 59–71.
- 73 Müller R, Stummann BM, Sisler EC, Serek M. Cultivar differences in regulation of ethylene production of miniature rose flowers (*Rosa hybrida* L.). *Gartenbauwissenschaft* 2001; **66**: 34–38.
- 74 Tan H, Liu X, Ma N, Xue J, Lu W et al. Ethylene-influenced flower opening and expression of genes encoding ETRs, CTRs, and EIN3 in two cut rose cultivars. *Postharvest Biol Technol* 2006; **40**: 97–105.
- 75 Xue J, Li Y, Tan H, Yang F, Ma N et al. Expression of ethylene biosynthetic and receptor genes in rose floral tissues during ethylene-enhanced flower opening. *J Exp Bot* 2008; **59**: 2161–2169.
- 76 Macnish AJ, Leonard RT, Borda AM, Nell TA. Genotypic variation in the postharvest performance and ethylene sensitivity of cut rose flowers. *J Hortic Sci* 2010; **45**: 790–796.
- 77 Clark DG, Dervinis C, Barrett JE, Nell TA. Using a seedling hypocotyl elongation assay as a genetic screen for ethylene sensitivity of seedling geranium cultivars. *Hort Technol* 2001; **11**: 297–302.
- 78 Kim HJ, Craig R, Brown KM. Genetic variation in ethylene responsiveness of *Regal Pelargonium*. *J Am Soc Hortic* 2006; **131**: 122–126.
- 79 Howard NP, Stimart D, de Leon N, Havey MJ, Martin W. Diallel analysis of floral longevity in *Impatiens walleriana*. *J Am Soc Hortic* 2012; **137**: 47–50.
- 80 Martin WJ, Stimart DP. Early generation evaluation in *Antirrhinum majus* for prediction of cut flower postharvest longevity. *J Am Soc Hortic* 2003; **128**: 876–880.
- 81 Weber JA, Martin WJ, Stimart DP. Genetics of postharvest longevity and quality traits in late generation crosses of *Antirrhinum majus* L. *J Am Soc Hortic* 2005; **130**: 694–699.
- 82 Martin WJ, Stimart DP. Genetic analysis of advanced populations in *Antirrhinum majus* L. with special reference to cut flower postharvest longevity. *J Am Soc Hortic* 2005; **130**: 434–441.
- 83 Friedman H, Hagiladi A, Resnick N, Barak A, Umiel N. Ethylene-insensitive related phenotypes exist naturally in a genetically variable population of *Dianthus barbatus*. *Theor Appl Genet* 2001; **103**: 282–287.
- 84 Krahel KH, Randle WM. Genetics of floral longevity in *Petunia*. *J Hortic Sci* 1999; **34**: 339–340.
- 85 Sun Y, Christensen B, Liu F, Wang H, Müller R. Effects of ethylene and 1-MCP (1-methylcyclopropene) on bud and flower drop in mini *Phalaenopsis* cultivars. *Plant Growth Regul* 2009; **59**: 83–91.
- 86 Chang YCA, Lin WL, Hou JY, Yen WY, Lee N. Concentration of 1-methylcyclopropene and the duration of its application affect anti-ethylene protection in *Phalaenopsis*. *Sci Hortic* 2013; **153**: 117–123.
- 87 Serek M, Reid MS. Ethylene and postharvest performance of potted *Kalanchoë*. *Postharvest Biol Technol* 2000; **18**: 43–48.
- 88 Macnish AJ, Irving DE, Joyce DC, Vithanage V, Wearing AH et al. Variation in ethylene-induced postharvest flower abscission responses among *Chamelacium* Desf. (*Myrtaceae*) genotypes. *Sci Hortic* 2004; **102**: 415–432.
- 89 Bicknell R. Breeding cut flower cultivars of *Leptospermum* using interspecific hybridisation. *New Zeal J Crop Hortic* 1995; **23**: 412–421.
- 90 Beal PR, Joyce A. Cut flower characteristics of terminal flowering tropical *Grevillea*: A brief review. *Aust J Exp Agric* 1999; **39**: 781–794.
- 91 Shibata M. Importance of genetic transformation in ornamental plant breeding. *Plant Biotech J* 2008; **25**: 3–8.
- 92 Riek JD, Debener T. Present use of molecular markers in ornamental breeding. In: Proc. 23rd Intl. Eucarpia Symp. (Sec. Ornamentals) on "Colourful Breeding and Genetics" Part II. Eds.: van Tuyl, J. M and de Vries, D. P. *Acta Hort* 2010; **855**: 77–84.
- 93 Yagi M. Application of DNA markers for breeding carnations resistant to bacterial wilt. *Jpn Agr Res Q* 2013; **47**: 29–35.
- 94 Izumikawa Y, Nakamura I, Mii M. Interspecific hybridization between *Kalanchoe blossfeldiana* and several wild *Kalanchoe* species with ornamental value. *Acta Hortic* 2009; **743**: 59–66.
- 95 Sasaki K, Aida R, Niki T, Yamaguchi H, Narumi T et al. High-efficiency improvement of transgenic *torenia* flowers by ion beam irradiation. *Plant Biotech J* 2008; **25**: 81–89.
- 96 Bologna P, Soto S, Prina AR, Borja M. X-rays as a tool for inducing variants of *Calibrachoa*. *Acta Hortic* 2012; **937**: 941–946.
- 97 McCallum CM, Comai L, Greene EA, Henikoff S. Targeting Induced Local Lesions In Genomes (TILLING) for plant functional genomics. *Plant Physiol* 2000; **123**: 439–442.



- 98 Dahmani-Mardas F, Troadec C, Boualem A, Lévêque S, Alsadon AA et al. Engineering melon plants with improved fruit shelf life using the TILLING Approach. *PLoS One* 2010; **5**: e15776.
- 99 Jiang P, Chen Y, Wilde HD. Optimization of EMS Mutagenesis on petunia for Tilling. *Adv Crop Sci Technol* 2014; **2**:141.
- 100 Xiong JS, Ding J, Li Y. Genome-editing technologies and their potential application in horticultural crop breeding. *Hort Res* 2015; **2**: 15019.
- 101 Savin KW, Baudinette SC, Graham MW, Michael MZ, Nugent GD et al. Antisense ACC Oxidase RNA delays carnation petal senescence. *J Hortic Sci* 1995; **30**: 970–972.
- 102 Kiss E, Veres A, Galli Z. Production of transgenic carnation with anti-sense ACS (1-aminocyclopropane-1-carboxylate synthase) gene. *Int J Hortic Sci* 2000; **6**: 103–107.
- 103 Iwazaki Y, Kosugi Y, Waki K, Yoshioka T, Satoh S. Generation and ethylene production of transgenic carnations harboring ACC synthase cDNA in sense or antisense orientation. *J Appl Hortic* 2004; **6**: 67–71.
- 104 Aida R, Yoshida T, Ichimura K, Goto R, Shibata M. Extension of flower longevity in transgenic torenia plants incorporating ACC oxidase transgene. *Plant Sci* 1998; **138**: 91–101.
- 105 Chen JC, Jiang CZ, Gookin TE, Hunter DA, Clark DG et al. Chalcone synthase as a reporter in virus-induced gene silencing studies of flower senescence. *Plant Mol Biol* 2004; **55**: 521–530.
- 106 Einset JW, Kopperud C. Antisense ethylene genes for begonia flowers. *Acta Hortic* 1995; **405**: 190–196.
- 107 Klee HJ, Hayford MB, Kretzmer KA, Barry GF, Kishore GM. Control of ethylene synthesis by expression of a bacterial enzyme in transgenic tomato plants. *Plant Cell* 1991; **3**: 1187–1193.
- 108 Wilkinson JQ, Lanahan MB, Clark DG, Bleecker AB, Chang C et al. A dominant mutant receptor from Arabidopsis confers ethylene insensitivity in heterologous plants. *Nat Biotechnol* 1997; **15**: 444–447.
- 109 Sanikhani M, Mibus H, Stummann BM, Serek M. *Kalanchoe blossfeldiana* plants expressing the Arabidopsis etr1-1 allele show reduced ethylene sensitivity. *Plant Cell Rep* 2008; **27**: 729–737.
- 110 Sriskandarajah S, Mibus H, Serek M. Transgenic *Campanula carpatica* plants with reduced ethylene sensitivity. *Plant Cell Rep* 2007; **26**: 805–813.
- 111 Sriskandarajah S, Mibus H, Serek M. Introduction of ethylene insensitivity in campanula carpatica flowers by Genetic Manipulation with Etr1-1. In: Proc. 6th International Postharvest Symposium. (eds.): Erkan, M. and Aksoy, U. *Acta Hort* 2010; **877**: 1111–1114.
- 112 Bovy AG, Angenent GC, Dons HJM, van Altvorst AN. Heterologous expression of the Arabidopsis etr1-1 allele inhibits the senescence of carnation flowers. *Mol Breeding* 1999; **5**: 301–308.
- 113 Gubrium EK, Clevenger DJ, Clark DG, Barrett JE, Nell TA. Reproduction and horticultural performance of transgenic ethylene-insensitive petunias. *J Am Soc Hortic* 2000; **125**: 277–281.
- 114 Cui ML, Takada K, Ma B, Ezura H. Overexpression of a mutated melon ethylene receptor gene Cm-ETR1/H69A confers reduced ethylene sensitivity in a heterologous plant, *Nemesia strumosa*. *Plant Science* 2004; **167**: 253–258.
- 115 Czarny JC, Grichko VP, Glick BR. Genetic modulation of ethylene biosynthesis and signaling in plants. *Biotechnol Adv* 2006; **24**: 410–419.
- 116 Gan S, Amasino RM. Inhibition of leaf senescence by autoregulated production of cytokinin. *Science New Series* 1995; **270**: 1986–1988.
- 117 Du L, Lou Q, Zhang X, Jiao S, Liu Y et al. Construction of flower-specific chimeric promoters and analysis of their activities in transgenic torenia. *Plant Mol Biol Rep* 2014; **32**: 234–245.
- 118 Wang H, Stier G, Lin J, Liu G, Zhang Z et al. Transcriptome changes associated with delayed flower senescence on transgenic petunia by inducing expression of etr1-1, a Mutant ethylene receptor. *PLoS One* 2013; **8**: e65800.
- 119 Caddick MX, Greenland AJ, Jepson I, Krause KP, Qu N et al. An ethanol inducible gene switch for plants used to manipulate carbon metabolism. *Nat Biotechnol* 1998; **16**: 177–180.
- 120 Garoosi GA, Salter MG, Caddick MX, Tomsett AB. Characterization of the ethanol-inducible alc gene expression system in tomato. *J Exp Bot* 2005; **56**: 1635–1642.
- 121 Moore I, Samalova M, Kurup S. Transactivated and chemically inducible gene expression in plants. *Plant J* 2006; **45**: 651–683.
- 122 Cobb D, Schneider N, Guo S, Humiston GA, Harriman B et al. Flower specific expression of an ethylene receptor (ETR1-1) confers ethylene insensitivity in transgenic Petunia. In: XXVth International Horticultural Congress Book of Abstracts, Toronto, 2002, 83.
- 123 Shaw JF, Chen HH, Tsai MF, Kuo CI, Huang LI. Extended flower longevity of petunia hybrida plants transformed with boers, a mutated ERS gene of *Brassica oleracea*. *Mol Breeding* 2002; **9**: 211–216.
- 124 Nukui H, Kudo S, Yamashita A, Satoh S. Repressed ethylene production in the gynoecium of long-lasting flowers of the carnation 'White Candle': Role of the gynoecium in carnation flower senescence. *J Exp Bot* 2004; **55**: 641–650.
- 125 Raffaeiner B, Serek M, Winkelmann T. *Agrobacterium tumefaciens*-mediated transformation of Oncidium and Odontoglossum orchid species with the ethylene receptor mutant gene etr1-1. *Plant Cell Tiss Org* 2009; **98**: 125–134.
- 126 Ciardi J, Barry K, Shibuya K, Nourizadeh S, Ecker J et al. Increased flower longevity in petunia through manipulation of ethylene signaling genes. In: Proceedings of the NATO Advanced Research Workshop on Biology and Biotechnology of the Plant Hormone Ethylene III., Eds. Vendrell, M., Klee, H., Pech, J. C. and Romojaro, F., 2003; 370–372.
- 127 Shibuya K, Berry KG, Ciardi JA, Loucas HM, Underwood BA et al. The central role of PhEIN2 in ethylene responses throughout plant development in petunia. *Plant Physiol* 2004; **136**: 2900–2912.
- 128 Chang X, Donnelly L, Sun D, Rao J, Reid MS et al. A petunia homeodomain-leucine zipper protein, PhHD-Zip, Plays an important role in flower senescence. *PLoS One* 2014; **9**: e88320.
- 129 Lusser M, Parisi C, Plan D, Rodríguez-Cerezo E. Deployment of new biotechnologies in plant breeding. *Nat Biotechnol* 2012; **30**: 231–239.
- 130 Podevin N, Devos Y, Davies HV, Nielsen KM. Transgenic or not? No simple answer! New biotechnology-based plant breeding techniques and the regulatory landscape. *EMBO reports* 2012; **13**: 1057–1061.
- 131 Carroll D. Genome engineering with zinc-finger nucleases. *J Genetics* 2011; **188**: 773–782.
- 132 Townsend JA, Wright DA, Winfrey RJ, Fu F, Maeder ML et al. High-frequency modification of plant genes using engineered zinc-finger nucleases. *Nature* 2009; **459**: 442–445.
- 133 Lor VS, Starker CG, Voytas DF, Weiss D, Olszewski NE. Targeted mutagenesis of the tomato PROCERA gene using transcription activator-like effector nucleases. *Plant Physiol* 2014; **166**: 1288.
- 134 Bortesi L, Fischer R. The CRISPR/Cas9 system for plant genome editing and beyond. *Biotechnol Adv* 2015; **33**: 41–52.
- 135 Mor Y, Spiegelstein H, Halevy AH. Inhibition of ethylene biosynthesis in carnation petals by cytokinin. *Plant Physiol* 1983; **71**: 541–546.
- 136 Hoeberichts FA, van Doorn WG, Vorst O, Hall RD, van Wordragen MF. Sucrose prevents up-regulation of senescence-associated genes in carnation petals. *J Exp Bot* 2007; **58**: 2873–2885.
- 137 van Doorn WG, Woltering EJ. Physiology and molecular biology of petal senescence. *J Exp Bot* 2008; **59**: 453–480.
- 138 Taverner EA, Letham DS, Wang J, Cornish E. Inhibition of carnation petal in-rolling by growth retardants and cytokinins. *Aust J Plant Physiol* 2000; **27**: 357–362.
- 139 Chang H, Jones ML, Banowets GM, Clarke DG. Overproduction of cytokinins in petunia flowers transformed with PSAG12-IPT delays corolla senescence and decreases sensitivity to ethylene. *Plant Physiol* 2003; **132**: 539–544.
- 140 Zakizadeh H, Lütken H, Sriskandarajah S, Serek M, Müller R. Transformation of miniature potted rose (*Rosa hybrida* cv. Linda) with PSAG12-ipt gene delays leaf senescence and enhances resistance to exogenous ethylene. *Plant Cell Rep* 2013; **32**: 195–205.
- 141 Trivellini A, Ferrante A, Vernieri P, Serra G. Effects of abscisic acid on ethylene biosynthesis and perception in Hibiscus rosa-sinensis L. flower development. *J Exp Bot* 2011; **62**: 5437–5452.
- 142 Nomura Y, Harada T, Morita S, Koshioka M et al. Role of ABA in triggering ethylene production in the gynoecium of senescing carnation flowers: Changes in ABA content and expression of genes for ABA biosynthesis and action. *Jpn Soc Hortic Sci* 2013; **82**: 242–254.
- 143 Liao WB, Zhang ML, Yu JH. Role of nitric oxide in delaying senescence of cut rose flowers and its interaction with ethylene. *Sci Hortic* 2013; **155**: 30–38.
- 144 Zhang C, Wang Y, Fu J, Dong L, Gao S et al. Transcriptomic analysis of cut tree peony with glucose supply using the RNA-Seq technique. *Plant Cell Rep* 2014; **33**: 111–129.
- 145 Müller R, Owen CA, Xu ZT, Welander M, Stummann BM. Genetic control of ethylene perception and signal transduction related to flower senescence. *J Food Agric Environ* 2003; **1**: 87–94.
- 146 van Doorn WG. Is petal senescence due to sugar starvation? *Plant Physiol* 2004; **134**: 35–42.
- 147 Shimizu H, Ichimura K. Effects of silver thiosulfate complex STS, sucrose and their combination on the quality and vase life of cut Eustoma flowers. *J Jpn Soc Hortic Sci* 2005; **74**: 381–385.
- 148 Zhang C, Liu M, Fu J, Wang Y, Dong L. Exogenous sugars involvement in senescence and ethylene production of tree peony 'Luoyang Hong' cut flowers. *Korean J Hortic Sci & Technol* 2012; **30**: 718–724.
- 149 Chen, MK, Hsu WH, Lee PF, Thiruvengadam M, Chen HI et al. The MADS box gene, FOREVER YOUNG FLOWER, acts as a repressor controlling floral organ senescence and abscission in Arabidopsis. *Plant J* 2011; **68**: 168–185.

- 150 Fernandez DE, Heck GR, Perry SE, Patterson SE, Bleecker AB *et al*. The embryo MADS domain factor AGL15 Acts post embryonically: Inhibition of perianth senescence and abscission via constitutive expression. *Plant Cell* 2000; **12**: 183–197.
- 151 Fang SC, Fernandez DE. Effect of regulated overexpression of the MADS domain factor AGL15 on flower senescence and fruit maturation. *Plant Physiol* 2002; **130**: 78–89.
- 152 Zheng Q, Zheng Y, Perry SE. AGAMOUS-Like15 promotes somatic embryogenesis in arabidopsis and soybean in part by the control of ethylene biosynthesis and response. *Plant Physiol* 2013; **161**: 2113–2127.
- 153 Fernandez DE, Wang CT, Zheng Y, Adamczyk BJ, Singhal R *et al*. The MADS-domain factors AGAMOUS-LIKE15 and AGAMOUS-LIKE18, along with SHORT VEGETATIVE PHASE and AGAMOUS-LIKE24, are necessary to block floral gene expression during the vegetative phase. *Plant Physiol* 2014; **165**: 1591–1603.
- 154 Tsai WC, Lee PF, Chen HI, Hsiao YY, Wei WJ *et al*. PeMADS6, a GLOBOSA/PISTILLATA-like Gene in *Phalaenopsis equestris* Involved in petaloid formation, and correlated with flower longevity and ovary development. *Plant Cell Physiol* 2005; **46**: 1125–1139.
- 155 Kandasamy MK, Deal RB, McKinney EC, Meagher RB. Silencing the nuclear actin-related protein AtARP4 in Arabidopsis has multiple effects on plant development, including early flowering and delayed floral senescence. *Plant J* 2005; **41**: 845–858.
- 156 Xu X, Jiang CZ, Donnelly L, Reid MS. Functional analysis of a RING domain ankyrin repeat protein that is highly expressed during flower senescence. *J Exp Bot* 2007; **58**: 3623–3630.
- 157 Jones RB, Serek M, Kuo CL, Reid MS. The effect of protein synthesis inhibition on petal senescence in cut bulb flowers. *J Am Soc Hortic Sci* 1994; **119**: 1243–1247.
- 158 Stier GN, Kumar P, Jiang CZ, Reid MS. Silencing of a proteasome component delays floral senescence. *Hortic Sci* 2010; **45**: S147.
- 159 Christensen B, Müller R. Improved postharvest life and ethylene tolerance in *Kalanchoe blossfeldiana* Transformed with rol-Genes of *Agrobacterium rhizogenes*. Proc. IXth Intl. Symp. on Postharvest Quality of Ornamental Plants Ed.: Ottosen, C.-O.. *Acta Hort*, 2009; **847**: 87–93.
- 160 Lütken H, Jensen EB, Christensen B. Development and evaluation of a non-GMO breeding technique exemplified by *Kalanchoë*. In: Proc. 7th, IS on In Vitro Culture and Horticultural Breeding, Ed.: D. Geelen. *Acta Hort* 2012; **961**: 51–58.
- 161 Lütken H, Wallström SV, Jensen EB, R. Inheritance of rol-genes from *Agrobacterium rhizogenes* through two generations in *Kalanchoë*. *Euphytica* 2012; **188**: 397–407.
- 162 da Silva JAT, Chin DP, Van PT, Mii M. Transgenic orchids. *Sci Hortic* 2011; **130**: 673–680.
- 163 Brand MH. Ornamental plant transformation. *J Crop Imp* 2006; **17**: 27–50.
- 164 Chandler SF, Sanchez C. Genetic modification; the development of transgenic ornamental plant varieties. *Plant Biotech J* 2012; **10**: 891–903.
- 165 Lusser M, Davies HV. Comparative regulatory approaches for groups of new plant breeding techniques. *N Biotechnol* 2013; **30**: 437–446.
- 166 Dirks R, van Dun K, de Snoo CB, van den Berg M, Lelivelt CL *et al*. Reverse breeding: A novel breeding approach based on engineered meiosis. *Plant Biotech J* 2009; **7**: 837–845.
- 167 Seeds of Change. *Nature* 2015; **520**: 131–132.
- 168 Cibus. Press Release, BASF and Cibus announce collaboration for herbicide tolerant crops. 2015. Available at [http://www.cibus.com/press\\_release.php?date=071007](http://www.cibus.com/press_release.php?date=071007) (accessed 20 April 2015).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 Unported License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>